

STATISTICAL PROPERTIES OF COUPLED EXPANDING MAPS ON A LATTICE WITH GENERAL INFINITE RANGE COUPLINGS AND HÖLDER DENSITIES

CHINMAYA GUPTA AND NICOLAI HAYDN

ABSTRACT. We continue the development of transfer operator techniques for expanding maps on a lattice coupled by general interaction functions. We obtain a spectral gap for an appropriately defined transfer operator, and, as corollaries, the existence of an invariant conformal probability measure for the system, exponential decay of correlations, the central limit theorem and the almost sure invariance principle.

1. INTRODUCTION

For one dimensional dynamical systems, the conditions under which there exists a unique ergodic invariant probability measure, supported on an invariant attractor and governing the dynamics of initial conditions in the basin of the attractor, are well understood. The complexity of the geometry in higher dimensions makes the problem much harder, however, in certain cases, significant results have been proven. For instance, the uniformly hyperbolic case has been presented in [Bow08], the non-uniformly hyperbolic case of the Hénon map in [BC91, BY93], and recently, the n dimensional analogue in [WY08]. In all, even though the 1 dimensional case is better understood in general than the n dimensional setting for $n > 1$, the theory of SRB measures for finite dimensional dynamical systems is fairly complete (for an excellent, though somewhat dated, overview, see [You02] and the references therein).

However, outside of a few settings, not much is known in the infinite dimensional setup. The primary problem is one of methodology, and the settings in which something can be said about an infinite dimensional dynamical system are those in which the techniques applicable in the finite dimensional case can be extended to infinite dimensions. Coupled maps on a lattice (CML) provide an example of infinite dimensional systems that can be studied by extensions of finite dimensional techniques, and for this reason, they have been extensively studied.

An excellent review of the theory of CML is provided in [Bun97]. In recalling the background on CML theory, we will be more restrictive; in particular, we will focus only on the development of ideas relevant to the results in this paper. An imprecise definition of a CML is provided in this section to set the context of the results that we mention. A precise formulation follows in later sections. Suppose we have a map τ on $[0, 1)$ and let $\Omega = [0, 1)^{\mathbb{Z}}$. We have an extended map $\bar{\tau}$ on the space Ω which can be defined as $(\bar{\tau}(x))_i = \tau(x_i)$ where $x \in \Omega$ and $x_i \in [0, 1)$. Suppose now we have interactions E between the different dynamical systems $(\tau_i, [0, 1))$. In the simplest case of E being the nearest neighbor diffusive coupling, we can specify the interactions as, for a given $0 < \epsilon \ll 1$,

$$(E(x))_i = (1 - \epsilon)x_i + \frac{\epsilon}{2}x_{i-1} + \frac{\epsilon}{2}x_{i+1}.$$

ϵ therefore becomes a parameter that tunes the strength of the interactions between the nodes of the lattice. We note also that this coupling specified has range 1, because only the closest neighbors to each node influence the state of that node. The system under study is now iterations of $E \circ \bar{\tau}$.

2000 *Mathematics Subject Classification.* 37A05, 37A25, 37A60, 60G10, 60F05, 80C20.

Key words and phrases. Coupled Lattice Maps, Statistical Properties of Dynamical Systems, Almost Sure Invariance Principle.

Bunimovich and Sinai [BS88] considered τ as expanding maps of the interval with nearest neighbor diffusive coupling with a non-constant diffusion strength, chosen so that the coupling map was onto. They established that if the interval map exhibits sufficient expansion, then there exists a unique invariant measure, mixing in time and space. They construct the invariant measure as the limit of the Gibbs measures for the finite dimensional projections of their coupled system. Then, in [PT92] the authors implement a numerical algorithm to extract orbits periodic in time and space for a 1-d lattice of Hénon maps, coupled by a weak nearest neighbor diffusive coupling. They define a new family of Lyapunov exponents that estimate the growth rate of spatially inhomogeneous perturbations.

Following the results in [BS88] and [PT92], a natural question arose: if the coupling is viewed as perturbation of the uncoupled system, what can happen if the coupling strength becomes large relative to the inherent stability of the uncoupled system? This question was studied in [BLMMR92], where the authors considered CML, where the maps on each node were one dimensional with a globally attracting stable periodic trajectory. These maps are coupled by a diffusive nearest neighbor coupling. The authors prove that if the coupling strength is sufficient large, then the phase space of the coupled map lattice is split into a complicated partition with many basins of different attractors.

The question of phase transitions for a one parameter family of finite-dimensional CML, with the coupling strength ϵ as the parameter of interest, was studied further, theoretically and numerically, by [KKN92]. They obtained sufficient conditions in terms of ϵ that the lattice have a continuum of ergodic components when the individual node maps are the doubling maps. They also produced an example of a function f for which the uncoupled system is mixing, whereas for a suitable ϵ , there are several domains in the phase space, interchanging with the period 2.

Keller and Künzle [KK92] then investigated transfer operator techniques for CML maps. They also considered expanding maps on each node of the lattice, and coupled them by a weak coupling (small ϵ). The authors established the spectral theory using spaces of bounded variation, closely following the setting in [BS88]. Then, in a three-part paper, Gundlach and Rand [GR93a, GR93b, GR93c] developed the stable manifold theory for coupled lattice maps with short or finite range interactions (Part I), and use this to establish the existence of a natural spatio-temporal measure that plays the role of the usual SRB measure in the case of temporal systems (Part III) and the existence and uniqueness of Gibbs states for higher dimensional symbolic systems by using thermodynamic formalism (Part II). The transfer operator approach of [KK92] was developed further by [Rug02] where the author established a spectral gap for weakly coupled real-analytic circle maps. In [Bar02], the author then modified the approach in [Rug02] to construct generalized transfer operators associated to potentials and establish a spectral gap for small potentials for weakly coupled analytic maps. Some limit theorems, such as the central limit theorem, moderate deviations, and a partial large deviations result were also established.

Returning to the setting of expanding real maps, [KL04] gave a proof of the existence, uniqueness and exponential mixing of invariant measures for weakly coupled lattice maps without cluster expansion. The coupling considered was finite range and had a very specific form, and the transfer operator was considered on the space of measures of bounded variation with absolutely continuous finite dimensional marginals. Subsequently, efforts were made to admit more general couplings and more general maps on the lattice nodes, and in [KL05] the authors study a one-dimensional lattice of weakly coupled piecewise expanding maps of the interval. Strong assumptions are still required on the coupling, however, the authors do not require that the coupling be a homeomorphism of the infinite dimensional state space. They prove that the transfer operator defined on an appropriate space of densities with bounded variations, with absolutely continuous finite dimensional marginals (with respect to the Lebesgue measure), has a spectral gap. This implies that there exists a measure with exponential decay of correlations in time and space. In [KL06] the authors extend the results

established in [KL05] to include lattices of any dimensions and couplings of infinite range with the coupling strength decaying exponentially in space.

In the setting of [KL04] and [KL06] the Bardet, Gouëzel and Keller [BGK07] prove the central limit theorem and the local limit theorem for Lipschitz functions depending on finitely many coordinates. The proof of the local limit theorem requires the additional assumption that for any compact interval J

$$\sigma\sqrt{2\pi n\mu_\epsilon} \left\{ x : \sum_{k=0}^{n-1} f \circ T_\epsilon^k(x) \in J \right\} \rightarrow |J|.$$

σ here is the variance in the central limit theorem. As in [KL04, KL06], the transfer operator is defined on the Banach space of measures of bounded variation with absolutely continuous finite dimensional marginals.

In this paper we continue the development of transfer operator techniques for studying existence and uniqueness of invariant measures corresponding to an initial potential in an appropriate class of potentials for expanding maps on a lattice coupled by an infinite range coupling. As in the setting of [KL04, KL06], we obtain a spectral gap for the transfer operator in an appropriate space of densities. In contrast to the setting of [KL04, KL06], we obtain our potentials and densities from a class of Hölder functions, and admit spatial couplings more general than those previously considered. We do not assume the existence of a reference measure; the theory is developed with respect to suitable potentials. Finally, we only require that the minimal expansion $\eta^{-1} > 1$ and the coupling strength $\epsilon > 0$ be related as $\epsilon\eta < 1$ for the existence, and uniqueness, of the invariant probability measure. We then use an abstract result of Gouëzel to obtain the almost sure invariance principle for our system. We also show that the invariant probability measure ν is conformal on open sets.

2. MAIN RESULTS.

In this section, we will list the main theorems that we prove. The precise definitions for the objects that appear here will follow in the body of the paper.

We start with a result known in the dynamical systems literature as “decay of correlations”. We establish this result by showing that an appropriate transfer operator \mathcal{L} has a spectral gap. For more details on the operator \mathcal{L} see sections 3.5, 4 and 5. We also check that the Lasota-Yorke inequality is stable under perturbation and in doing so, establish the almost sure invariance principle by using [Gou10]. We use the space \mathcal{C} of Hölder continuous functions with the norm $\|\cdot\| = |\cdot|_\infty + |\cdot|_\beta$ where $|\cdot|_\beta$ is the β -Hölder constant (the precise definition is below).

Theorem 2.1. *Let T be the coupled lattice map on the lattice system Ω and \mathcal{C} the space of Hölder continuous functions on Ω .*

Then for any potential function $f \in \mathcal{C}$ there exists an invariant measure ν which is g -conformal where $g = f + h - f \circ T$ for an $h \in \mathcal{C}$. Moreover there exists a constant $0 < \varsigma < 1$ with the property that for any $\Phi_1, \Phi_2 \in \mathcal{C}$ there exists a constant C_1 such that

$$\left| \int \Phi_1 \circ T^n \Phi_2 d\nu - \int \Phi_1 d\nu \int \Phi_2 d\nu \right| \leq C_1 |\Phi_1|_\infty \|\Phi_2\| \varsigma^n.$$

Finally, by using an abstract result from [Gou10], and Theorem 4.1, we prove that

Theorem 2.2. *The system satisfies the almost sure invariance principle.*

Various statistical properties such as the law of iterated logarithms, the weak invariance principle and the central limit theorem now follow as corollaries.

Corollary 2.3. (CLT) *For observables $\Phi \in \mathcal{C}$ one has*

$$\mathbb{P} \left(\frac{\sum_{j=0}^{n-1} \Phi \circ T^j - n\nu(\Phi)}{\sigma\sqrt{n}} \leq t \right) \rightarrow N(t)$$

as $n \rightarrow \infty$ where T is the coupled lattice map and $N(t)$ is the normal distribution.

A technical result along the way is the following proposition:

Proposition 2.4. *Let $\Phi \in \mathcal{C}$. Then $\mathcal{L}(\Phi) \in \mathcal{C}$ and moreover there exists a constant $\theta_1 \in (0, 1)$ and a constant $C_2 > 0$ such that*

$$|\mathcal{L}^n(\Phi)|_\beta \leq |\Phi|_\beta \theta_1^n + C_2 |\Phi|_\infty$$

The proposition is used to prove the existence of the invariant probability measure ν which satisfies $\mathcal{L}^*\nu = \nu$. This result is proved as Theorem 4.1. As a corollary to this theorem, with suitable modifications to the classical techniques we prove that the essential spectral radius of \mathcal{L} is strictly smaller than the spectral radius of \mathcal{L} (which is 1).

Finally, we make a note regarding constants. We denote ‘global’ constants by C_1, C_2, \dots throughout the paper and ‘local’ constants by c_1, c_2, \dots for each lemma or theorem. The constant C_E tunes the strength of interactions between the maps on the lattice.

3. THE UNCOUPLED SYSTEM.

First, we need to define what the admissible class of observables is. In order to study observables (and potentials) on an infinite lattice, we need to define how well approximated these observables are by restrictions to finite sub-lattices. Let $\Lambda = \mathbb{Z}$ and $\Omega = I^{\mathbb{Z}}$ the lattice space where I is the unit interval. Define

$$\Lambda_k = (-k, -k+1, \dots, -1, 0, 1, \dots, k-1, k).$$

Assume the following:

3.1. Assumptions on the metric.

(d1) Let $d_I : I \rightarrow \mathbb{R}^+$ be a metric on the unit interval I and pick $\theta \in (0, 1)$. Define a metric d on Ω by

$$(1) \quad d(x, y) = \sup_{k \in \mathbb{Z}} \theta^{|k|} d_I(x_k, y_k).$$

where $x = (\dots, x_{-1}, x_0, x_1, x_2, \dots)$, $y = (\dots, y_{-1}, y_0, y_1, y_2, \dots)$ are points in Ω .

3.2. Assumptions on the map on I . The global map on Ω is thought to be composed of individual maps on the nodes I . We list below the conditions that our map satisfies (we note that these are not the most general conditions under which our theorem can be stated, but in order to keep our reasoning transparent, we do not pursue a more general setting).

- (τ 1) The map $\tau : I \rightarrow I$ has full branches. As a consequence, for each $x \in I$, $b = \#\{\tau^{-1}x\}$ is constant. This also implies that the map τ has at least one fixed point. Denote one such fixed point by p_τ .
- (τ 2) The map $\tau : I \rightarrow I$ is expanding. This is taken to mean that there exists a constant $\eta \in (0, 1)$ such that for every inverse branch ζ of τ one has $d(\zeta_x, \zeta_y) \leq \eta d(x, y)$ (ζ_x, ζ_y are the images of x, y under ζ , i.e. $\tau\zeta_x = x, \tau\zeta_y = y$).

Define $i_k : \Omega_k \rightarrow \Lambda$ as $i_k(x) = \begin{cases} x_i & \text{if } |i| \leq k \\ p_\tau & \text{otherwise} \end{cases}$ and put $\pi_k : \Lambda \rightarrow \Lambda$ for the ‘projection’ which is given by $\pi_k x = i_k(x|_{\Omega_k})$ where $\Omega_k = I^{\Lambda_k}$.

For $\Phi \in C(\Omega)$ we define the ‘restriction’ to Ω_k as

$$\Phi_k = \Phi \circ \pi_k$$

that is $\Phi_k(x) = \Phi \circ i_k(x|_{\Omega_k})$. A consequence of the fact that outside of the lattice Ω_k we have chosen $\pi_k(x)$ to be equal to p_τ is that

$$\bar{\tau}^i \pi_k(x) = \pi_k \bar{\tau}^i(x).$$

3.3. Definition of the function space \mathcal{C} . For $\Phi \in C(\Omega)$ we define the Hölder constant

$$|\Phi|_\beta := \sup_{k \in \mathbb{N}} |\Phi_k|_\beta$$

where

$$|\Phi_k|_\beta := \sup_{x, y \in \Omega} \frac{|\Phi_k(x) - \Phi_k(y)|}{d(x, y)^\beta}.$$

Then

$$\|\cdot\| = |\cdot|_\infty + |\cdot|_\beta$$

defines a norm and we define the function space

$$\mathcal{C}_\beta = \{\Phi \in C(\Omega) : \|\Phi\| < \infty\}.$$

We will also sometimes need the variation semi-norm for some $\alpha \in (0, 1)$ given by

$$V_\alpha(\Phi) := \sup_{k \in \mathbb{N}} \frac{\text{var}_k(\Phi)}{\alpha^k}$$

where

$$\text{var}_k(\Phi) = \sup_{x \in \Omega} |\Phi(x) - \Phi_k(x)|$$

is the k th variation of Φ . If $\alpha \geq \theta^\beta$ then $V_\alpha(\Phi) \leq |\Phi|_\beta$ for all $\Phi \in \mathcal{C}_\beta$. Therefore, in what follows, we fix some $\alpha \geq \theta^\beta$ and, instead of writing \mathcal{C}_β , we only write \mathcal{C} .

3.4. Defining the positive operators P and L . Let $f \in \mathcal{C}$ be a potential function and define for the finite sub-lattice Λ_k the transfer operator P_k for $\bar{\tau}|_{\Omega_k}$ by

$$P_k(\Phi)(x) = \frac{1}{b_k} \sum_{|\zeta|=k} \exp(f(i_k \zeta_x)) \Phi(i_k \zeta_x)$$

$x \in \Omega$, where the summation is over inverse branches ζ of $\bar{\tau}|_{\Omega_k}$, i.e. $\bar{\tau} \circ \zeta$ is the identity and moreover ζ_x the image of x under ζ has the property that $(\zeta_x)_j = x_j$ for $|j| > k$. For the normalising factor one has $b_k = \#\{\bar{\tau}^{-1}x|_{\Omega_k}\} = b^{2k+1}$ as $\#\{\bar{\tau}^{-1}x\} = b$.

Clearly, P_k is a well defined, positive, bounded linear operator on functions on Ω . The following lemma serves to define the transfer operator P by taking a limit k to infinity.

Lemma 3.1. *Let $f : \Omega \rightarrow \mathbb{R}$ such that $|f|_\infty + V_\alpha(f) < \infty$. Then*

- (I) P_k is uniformly (in k) bounded.
- (II) For Φ such that $|\Phi|_\infty + V_\alpha(\Phi) < \infty$ the sequence $(P_k(\Phi))$ is Cauchy for each $x \in \Omega$.
- (III) $V_\alpha(P_k \Phi) \leq C_3 |P \Phi|_\infty \quad \forall k$ for some constant C_3 .

Proof. (I) Clearly $|P_k \Phi|_\infty \leq e^{|f|_\infty} |\Phi|_\infty$, which implies P_k is bounded uniformly in k .

(II) For $k_1 < k_2$ one has

$$|P_{k_2}(\Phi)(x) - P_{k_1}(\Phi)(x)| = \left| \sum_{|\zeta'|=k_2} \frac{\exp(f \zeta'_x) \Phi(\zeta'_x)}{b_{k_2}} - \sum_{|\zeta|=k_1} \frac{\exp(f \zeta_x) \Phi(\zeta_x)}{b_{k_1}} \right|$$

where ζ' are inverse branches in $\bar{\tau}^{-1}|_{\Omega_{k_2}}$ and ζ are inverse branches in $\bar{\tau}^{-1}|_{\Omega_{k_1}}$.

The lattice Λ_{k_2} contains $2(k_2 - k_1)$ elements more than Λ_{k_1} , and so $b_{k_2} = b^{2(k_2 - k_1)} b_{k_1}$. Therefore the above sum simplifies as

$$\begin{aligned}
 & \left| \sum_{|\zeta'|=k_2} \frac{\exp(f\zeta'_x)\Phi(\zeta'_x)}{b_{k_2}} - \sum_{|\zeta|=k_1} \frac{\exp(f\zeta_x)\Phi(\zeta_x)}{b_{k_1}} \right| \\
 & \leq \sum_{|\zeta|=k_1} \left[\frac{\left(\sum_{\zeta'|\Omega_{k_1}=\zeta} \exp(f\zeta'_x)\Phi(\zeta'_x) \right)}{b_{k_2}} - \frac{\exp(f\zeta_x)\Phi(\zeta_x)}{b_{k_1}} \right] \\
 (2) \quad & = \sum_{|\zeta|=k_1} \frac{\exp(f\zeta_x)\Phi(\zeta_x)}{b^{2k_1+1}} \left[\left(\sum_{\zeta'|\Omega_{k_1}=\zeta} \exp(f\zeta'_x - f\zeta_x) \frac{\Phi(\zeta'_x)}{\Phi(\zeta_x)} \right) \frac{b^{2k_1+1}}{b^{2k_2+1}} - 1 \right]
 \end{aligned}$$

Since $\exp(f\zeta'_x - f\zeta_x) \leq e^{V_\alpha(f)} \leq 1 + V_\alpha(f)\alpha^{k_1} + o(\alpha^{k_1})$ and

$$\left| \frac{\Phi(\zeta'_x)}{\Phi(\zeta_x)} - 1 \right| \leq \Phi(\zeta_x) V_\alpha(\Phi) \alpha^{k_1} \leq |\Phi|_\infty V_\alpha(\Phi) \alpha^{k_1}.$$

Therefore (for some c_1)

$$\left(\sum_{\zeta'|\Omega_{k_1}=\zeta} \exp(f\zeta'_x - f\zeta_x) \frac{\Phi(\zeta'_x)}{\Phi(\zeta_x)} \right) = \sum_{\zeta'|\Omega_{k_1}=\zeta} \left(1 + c_1 \alpha^{k_1} + o(\alpha^{k_1}) \right) = b^{2k_2-2k_1} (1 + c_1 \alpha^{k_1} + o(\alpha^{k_1}))$$

from where it follows that

$$\left[\left(\sum_{\zeta'|\Omega_{k_1}=\zeta_x} \exp(f\zeta'_x - f\zeta_x) \frac{\Phi(\zeta'_x)}{\Phi(\zeta_x)} \right) \frac{b^{2k_1+1}}{b^{2k_2+1}} - 1 \right] \leq c_1 \alpha^{k_1} + o(\alpha^{k_1}).$$

Therefore,

$$|P_{k_2}(\Phi)(x) - P_{k_1}(\Phi)(x)| \leq c_2 |P_{k_1}\Phi|_\infty \alpha^{k_1} \leq c_3 \alpha^k$$

for some constants c_2, c_3 as by part (I) $|P_{k_1}\Phi|_\infty$ is uniformly bounded.

(III) This follows from the first inequality in the last line of estimates. \square

Since by Lemma 3.1, for each $x \in \Omega$, the sequence $P_k(\Phi)(x)$ is a Cauchy sequence (of real numbers) we now define the operator P for the infinite lattice system as the pointwise limit:

$$P(\Phi)(x) = \lim_{k \rightarrow \infty} P_k(\Phi)(x).$$

Lemma 3.2. P is a non-negative and continuous operator on \mathcal{C} .

Proof. Clearly P is non-negative as the approximations P_k are non-negative. Since P is a linear operator, it is enough to show continuity at the origin. For $\Phi \in \mathcal{C}$ we see that P is a bounded operator in the $|\cdot|_\infty$ -norm as

$$|P(\Phi)| \leq e^{|f|_\infty} |\Phi|_\infty.$$

Now let $x, y \in \Omega$, then

$$\begin{aligned}
 |P\Phi(x) - P\Phi(y)| &\leq \lim_k \frac{1}{b_k} \sum_{|\zeta|=k} \left| e^{f(i_k \zeta_x)} \Phi(i_k \zeta_x) - e^{f(i_k \zeta_y)} \Phi(i_k \zeta_y) \right| \\
 &\leq \lim_k \frac{1}{b_k} \left(\sum_{\zeta} e^{f(i_k \zeta_x)} |\Phi(i_k \zeta_x) - \Phi(i_k \zeta_y)| + |\Phi|_{\infty} \sum_{\zeta} e^{f(i_k \zeta_y)} \left| e^{f(i_k \zeta_y) - f(i_k \zeta_x)} - 1 \right| \right) \\
 &\leq e^{|f|_{\infty}} |\Phi|_{\beta} d(x, y)^{\beta} + c_1 |\Phi|_{\infty} |f|_{\beta} d(x, y)^{\beta}
 \end{aligned}$$

for some c_1 as $d(i_k \zeta_x, i_k \zeta_y) \leq c_2 d(x, y)$ ($c_2 > 0$). Hence $|P\Phi|_{\beta} \leq c_3(|\Phi|_{\beta} + |\Phi|_{\infty}) = c_3 \|\Phi\|$ for some constant c_3 which is independent of Φ . \square

Note that the space Ω with the metric d is convex, compact and separable. Separability follows from the fact that for every $k \in \mathbb{N}$ there are finitely many points in $i_k \Omega_k$ that are ϑ^k -dense in Ω for any $\vartheta \in (0, 1)$. In this way one produces a countable dense set in Ω . This implies that the set \mathcal{M} of probability measures on Ω is compact in the weak* topology. Thus following [Bow08] we can define an operator $\mathcal{M} : \mathcal{M} \rightarrow \mathcal{M}$ by $\mathcal{M}\nu = \frac{P^*\nu}{P^*\nu(\mathbb{1})}$ where $\nu \in \mathcal{M}$. By the theorem of Schauder-Tychonoff \mathcal{M} has a fixed point ν in \mathcal{M} . Thus $P^*\nu = \lambda\nu$, where $\lambda = P^*\nu(\mathbb{1})$.

Definition 3.3. Let $f \in \mathcal{C}$ and put $B(z) = \exp\left(|f|_{\beta} \frac{\eta^{\beta}}{1-\eta^{\beta}} z^{\beta}\right)$, $z \geq 0$. Define the function set

$$\Delta_f := \{\Phi \in C(\Omega) : \Phi \geq 0, \nu(\Phi) = 1, \Phi(x) \leq B(d(x, y))\Phi(y) \forall x, y \in \Omega\}.$$

Notice that $B(\eta z) e^{|f|_{\beta} \eta^{\beta} z^{\beta}} = B(z)$.

Lemma 3.4. $\Delta_f \subset \mathcal{C}$.

Proof. For $\Phi \in \Delta_f$ one has $\Phi(x)/\Phi(y) \leq B(d(x, y))$ and $\Phi(y)/\Phi(x) \leq B(d(x, y))$ which implies

$$|\Phi(x) - \Phi(y)| = \left| \frac{\Phi(x)}{\Phi(y)} - 1 \right| |\Phi(y)| \leq |\Phi(y)| |B(d(x, y)) - 1| \leq |\Phi|_{\infty} \left[d(x, y)^{\beta} |f|_{\beta} \frac{\eta^{\beta}}{1-\eta^{\beta}} + o(d(x, y)^{\beta}) \right]$$

where $|\Phi|_{\infty} \leq B(1)$ as the diameter of Ω is equal to 1 and $\nu(\Phi) = 1$ (i.e. $\inf \Phi \leq 1$). This implies that $|\Phi|_{\beta} \leq |f|_{\beta} \left(\frac{\eta^{\beta}}{1-\eta^{\beta}} + c_1 \right) < \infty$ for some c_1 . Hence $\|\Phi\| < \infty$. \square

In order to apply the theorem of Schauder-Tychonoff we must first show that the operator $\frac{1}{\lambda}P$ maps Δ_f into itself.

Lemma 3.5. $\frac{1}{\lambda}P$ maps Δ_f into itself.

Proof. Clearly $\nu(\frac{1}{\lambda}P\Phi) = 1$ for all $\Phi \in \Delta_f$. Since P is a positive operator we also get $\frac{1}{\lambda}P\Phi \geq 0$ for all $\Phi \in \Delta_f$. It remains to verify the regularity property. Since

$$f(i_k \zeta_x) \leq f(i_k \zeta_y) + |f|_{\beta} d(i_k \zeta_x, i_k \zeta_y)^{\beta} \leq f(i_k \zeta_y) + |f|_{\beta} \eta^{\beta} d(x, y)^{\beta}$$

one obtains

$$\begin{aligned}
 \frac{1}{\lambda}P\Phi(x) &= \frac{1}{\lambda} \lim_{k \rightarrow \infty} \frac{1}{b_k} \sum_{|\zeta|=k} e^{f(i_k \zeta_x)} \Phi(i_k \zeta_x) \\
 &\leq \frac{1}{\lambda} \lim_{k \rightarrow \infty} \frac{1}{b_k} \sum_{|\zeta|=k} e^{f(i_k \zeta_y)} \Phi(i_k \zeta_y) B(\eta d(x, y)) e^{|f|_{\beta} \eta^{\beta} d(x, y)^{\beta}} \\
 &\leq \frac{1}{\lambda} P\Phi(y) B(\eta d(x, y)) e^{|f|_{\beta} \eta^{\beta} d(x, y)^{\beta}} \\
 &\leq \frac{1}{\lambda} P\Phi(y) B(d(x, y)).
 \end{aligned}$$

Therefore, $\frac{1}{\lambda}P\Phi \in \Delta_f$. □

Lemma 3.6. *There exist a unique $h \in \Delta_f$ so that $Ph = \lambda h$ and moreover h is strictly positive.*

Proof. The set Δ_f is convex and equicontinuous by Lemma 3.4. Therefore Δ_f is compact in the $|\cdot|_\infty$ -norm by Arzela-Ascoli and $\frac{1}{\lambda}P$ has by Schauder-Tychonoff a fixed point $h \in \Delta_f$. That is $Ph = \lambda h$. To see that h is strictly positive assume that h has a zero at $x \in \Omega$, i.e. $h(x) = 0$. Then

$$0 = \frac{1}{\lambda^n} P^n h(x) = \frac{1}{\lambda^n} \lim_k P_k^n h(x) = \frac{1}{\lambda^n} \lim_{k \rightarrow \infty} \frac{1}{b_k} \sum_{\zeta} e^{f^n(i_k \zeta_x)} h(i_k \zeta_x)$$

where the sum is over all inverse branches ζ of $\bar{\tau}^{-n}$ in $i_k \Lambda_k$. Since $h \geq 0$ this implies that $h(\zeta_x) = 0$ for all inverse branches ζ of $\bar{\tau}^n$. Since the set $\bigcup_n \bigcup_{\zeta \in \bar{\tau}^{-n}} \zeta_x$ is dense in Ω and h is continuous we conclude that h is identically zero which contradicts the assumption $\nu(h) = 1$.

To obtain uniqueness of h assume that there is a second eigenfunction $h' \in \Delta_f$ so that $Ph' = \lambda h'$ and put $t = \inf \frac{h'}{h}$. By convexity of Δ_f one has $h - th \in \Delta_f$ and by choice of t there exists an $x \in \Omega$ so that $(h - th')(t) = 0$. By the argument above we conclude that $h - th'$ must vanish identically, which is impossible. Hence h is unique. □

We now define the normalized transfer operator $L : \mathcal{C} \rightarrow \mathcal{C}$ by putting $L(\Phi) := P(h\Phi)/(\lambda h)$ for $\Phi \in \mathcal{C}$. Note that $L(\Phi)$ is well defined since $h > 0$ and has the potential function $g = f - \log \lambda - \log h \circ \bar{\tau} + \log h$. Moreover L has the (dominant) simple eigenvalue 1 with eigenfunction $\mathbb{1}$ as $L(\mathbb{1}) = \mathbb{1}$. The associated eigen-functional $\mu = h\nu$ is a $\bar{\tau}$ -invariant probability measure. Define, also, $L_k(\Phi)$ as follows:

$$L_k(\Phi)(x) = \frac{1}{b_k} \sum_{|\zeta|=k} e^{g(i_k \zeta_x)} \Phi(i_k \zeta_x)$$

where $g(x) = f(x) - \log \lambda - \log h \circ \bar{\tau}(x) + \log h(x)$. Note that by definition, $|g|_\beta \leq |f|_\beta + 2|h|_\beta < \infty$, and $|g|_\infty < \infty$.

Hence, we state a corollary to Lemma 3.1:

Corollary 3.7. *For each x , and for each Φ , the sequence $L_k(\Phi)(x)$ is Cauchy, and hence it converges to $L(\Phi)(x)$.*

Proof. The fact that $L_k(\Phi)(x)$ is Cauchy, and hence has a point wise limit, follows directly from Lemma 3.1. Also, notice that

$$\begin{aligned} L(\Phi)(x) &= \frac{P(h\Phi)(x)}{\lambda h(x)} \\ &= \lim_{k \rightarrow \infty} \frac{P_k(h\Phi)(x)}{\lambda h(x)} \\ &\geq \lim_{k \rightarrow \infty} \frac{1}{b_k} \sum_{|\zeta|=k} e^{f(i_k \zeta_x)} h(i_k \zeta_x) \Phi(i_k \zeta_x) \frac{1}{\lambda h(x)}. \end{aligned}$$

The term in the last summation can be written as

$$e^{f(i_k \zeta_x)} h(i_k \zeta_x) \Phi(i_k \zeta_x) \frac{1}{\lambda h(x)} = e^{g(i_k \zeta_x)} \Phi(i_k \zeta_x) \frac{h(\tau(i_k \zeta_x))}{h(x)},$$

and so, because $\text{var}(h) < \infty$, for any $\epsilon > 0$, we have that

$$L(\Phi)(x) \geq (1 - \epsilon) \lim_{k \rightarrow \infty} L_k(\Phi)(x).$$

Similarly, we obtain that $L(\Phi)(x) \leq (1 + \epsilon) \lim_{k \rightarrow \infty} L_k(\Phi)(x)$. This completes the proof. □

3.5. Lasota-Yorke inequality for L . We now establish a Lasota-Yorke inequality for the operator L . We do this by obtaining the corresponding inequality for each approximation L_k . We use the following notation: For a function Φ we denote by $\Phi^{(n)} = \sum_{j=0}^{n-1} \Phi \circ \bar{\tau}$ its n th ergodic sum. We shall need the following technical estimate.

Lemma 3.8. *Let $f \in \mathcal{C}$ be a potential and let $g = f - \log \lambda - \log h \circ \bar{\tau} + \log h$. Then for all x and $y \in \Omega$:*

$$\left| g^{(n)}(i_k \zeta_x) - g^{(n)}(i_k \zeta_y) \right| \leq \left(3|h|_\beta + \frac{\eta^\beta}{1 - \eta^\beta} |f|_\beta \right) d(x, y)^\beta.$$

Proof. Since

$$g^{(n)}(i_k \zeta_x) = \sum_{j=0}^{n-1} g \circ \bar{\tau}^j(i_k \zeta_x) = \sum_{j=0}^{n-1} f \circ \bar{\tau}^j(i_k \zeta_x) - n \log \lambda + h(i_k \zeta_x) - h \circ \bar{\tau}^n(i_k \zeta_x)$$

we get that

$$g^{(n)}(i_k \zeta_x) - g^{(n)}(i_k \zeta_y) = f^{(n)}(i_k \zeta_x) - f^{(n)}(i_k \zeta_y) + h(i_k \zeta_x) - h(i_k \zeta_y) + h \circ \bar{\tau}^n(i_k \zeta_y) - h \circ \bar{\tau}^n(i_k \zeta_x).$$

Note that $|h \circ \bar{\tau}^n(i_k \zeta_y) - h \circ \bar{\tau}^n(i_k \zeta_x)|$ is bounded by $|h|_\beta d(i_k x|_{\Lambda_k}, i_k y|_{\Lambda_k})^\beta \leq |h|_\beta d(x, y)^\beta$ and $|h(i_k \zeta_x) - h(i_k \zeta_y)|$ is bounded by $|h|_\beta \eta^{\beta n} d(x, y)^\beta$. Define $\tau_{(k)} := \bar{\tau} \circ i_k$. The above combined with

$$\begin{aligned} \left| f^{(n)}(i_k \zeta_x) - f^{(n)}(i_k \zeta_y) \right| &\leq \sum_{i=0}^{n-1} |f \circ \bar{\tau}^i(i_k \zeta_x) - f \circ \bar{\tau}^i(i_k \zeta_y)| \\ &\leq \sum_{i=0}^{n-1} \left| f(i_k(\tau_{(k)}^i \zeta_x)) - f(i_k(\tau_{(k)}^i \zeta_y)) \right| \\ &\leq \sum_{i=0}^{n-1} |f(i_k((\tau^i i_k \zeta_x))|_{\Omega_k}) - f(i_k((\tau^i i_k \zeta_y))|_{\Omega_k})| \\ &\leq |f|_\beta \sum_{i=0}^{n-1} d(i_k(\bar{\tau}_{(k)}^i \zeta_x), i_k(\bar{\tau}_{(k)}^i \zeta_y))^\beta \\ &\leq |f|_\beta d(x, y)^\beta \sum_{i=0}^{n-1} \eta^{\beta(n-i)}. \end{aligned}$$

proves the desired bound. □

Proposition 3.9 (Lasota-Yorke inequality for L_k). *Let $\Phi \in \mathcal{C}$ and f, h, τ be as before. Let $k \in \mathbb{Z}^+$. Then $\exists C_4 > 0$ (depending on f, h and τ) such that*

$$|L_k^n(\Phi)|_\beta \leq \left(|\Phi|_\beta \eta^{\beta n} + C_4 |\Phi|_\infty \right) L_k^n(\mathbf{1}).$$

Proof. For $x, y \in \Omega$:

$$\begin{aligned}
|L_k^n(\Phi)(x) - L_k^n(\Phi)(y)| &\leq \frac{1}{b_k} \sum_{|\zeta|=1} \left| e^{g^{(n)}(i_k \zeta_x)} \Phi(i_k \zeta_x) - e^{g^{(n)}(i_k \zeta_y)} \Phi(i_k \zeta_y) \right| \\
&\leq \frac{1}{b_k} \left[\sum_{\zeta} e^{g^{(n)}(i_k \zeta_x)} |\Phi(i_k \zeta_x) - \Phi(i_k \zeta_y)| + \sum_{\zeta} |\Phi|_{\infty} \left| e^{g^{(n)}(i_k \zeta_x)} - e^{g^{(n)}(i_k \zeta_y)} \right| \right] \\
&\leq \frac{1}{b_k} \left[\sum_{\zeta} |\Phi|_{\beta} \eta^{\beta n} d(x, y)^{\beta} L_k^n(\mathbb{1}) \right. \\
&\quad \left. + \sum_{\zeta} |\Phi|_{\infty} e^{g^{(n)}(i_k \zeta_y)} \left| 1 - \exp(g^{(n)}(i_k \zeta_x) - g^{(n)}(i_k \zeta_y)) \right| \right]
\end{aligned}$$

as $d(i_k \zeta_x, i_k \zeta_y) \leq \eta^n d(x, y)$. By Lemma 3.8, we have

$$\left| 1 - \exp(g^{(n)}(i_k \zeta_x) - g^{(n)}(i_k \zeta_y)) \right| \leq c_1 d(x, y)^{\beta} + o(d(x, y)^{\beta}) \leq c_2 d(x, y)^{\beta}$$

where $c_1 \leq 3|h|_{\beta} + \frac{\eta^{\beta}}{1-\eta^{\beta}}|f|_{\beta}$. Consequently

$$\sup_{x \neq y} \frac{|L_k^n(\Phi)(x) - L_k^n(\Phi)(y)|}{d(x, y)^{\beta}} \leq |\Phi|_{\beta} \eta^{\beta n} L_k^n(\mathbb{1}) + c_2 |\Phi|_{\infty} L_k^n(\mathbb{1}).$$

□

Now we can prove the 'Lasota-Yorke' inequality for the operator L for the uncoupled map $\bar{\tau}$ on the full, infinite lattice Ω .

Theorem 3.10. *Let $\Phi \in \mathcal{C}$ and let f, h, τ be as before, and $n \in \mathbb{N}$. There exists a constant $C_5 > 0$ depending only on f, h, τ such that*

$$|L^n(\Phi)|_{\beta} \leq |\Phi|_{\beta} \eta^{\beta n} + C_5 |\Phi|_{\infty}.$$

Proof. Recall η from Assumption ($\tau 2$). Let $k_2 \geq k_1 \in \mathbb{Z}^+$. Then for all $x, y \in \Omega$ one has

$$\begin{aligned}
|L_{k_2}^n(\Phi)(y) - L_{k_1}^n(\Phi)(x)| &\leq |L_{k_1}^n(\Phi)(x) - L_{k_1}^n(\Phi)(y)| + |L_{k_1}^n(\Phi)(y) - L_{k_2}^n(\Phi)(y)| \\
&\leq \left[|\Phi|_{\beta} \eta^{\beta n} + C_4 |\Phi|_{\infty} \right] L_{k_1}^n(\mathbb{1}) d(x, y)^{\beta} + C_3 |L_{k_1}^n(\Phi)|_{\infty} \alpha^{k_1} \\
&\leq \left[|\Phi|_{\beta} \eta^{\beta n} + C_4 |\Phi|_{\infty} \right] L_{k_1}^n(\mathbb{1}) d(x, y)^{\beta} + C_3 e^{|g|_{\infty}} |\Phi|_{\infty} \alpha^{k_1},
\end{aligned}$$

where the second line uses Lemmas 3.9 and 3.1 (III) while the last line follows from the proof of Lemma 3.1 (I). Letting $k_2 \rightarrow \infty$ we see that

$$|L^n(\Phi)(y) - L_{k_1}^n(\Phi)(x)| \leq \left[|\Phi|_{\beta} \eta^{\beta n} + C_4 |\Phi|_{\infty} \right] L_{k_1}^n(\mathbb{1}) d(x, y)^{\beta} + C_3 e^{|g|_{\infty}} |\Phi|_{\infty} \alpha^{k_1}$$

and then on letting $k_1 \rightarrow \infty$, and recalling that $L_{k_1}(\mathbb{1}) \rightarrow L(\mathbb{1}) = \mathbb{1}$ we get

$$|L^n(\Phi)(y) - L^n(\Phi)(x)| \leq \left[|\Phi|_{\beta} \eta^{\beta n} + C_4 |\Phi|_{\infty} \right] d(x, y)^{\beta}$$

The proof is complete on dividing by $d(x, y)^{\beta}$ and taking the supremum. We put $C_5 = C_4$. □

4. THE PERTURBATION E .

Denote by $\sigma : \Omega \rightarrow \Omega$ the shift map which is given by $\sigma((x_i)_{i=-\infty}^\infty) = (x_{i+1})_{i=-\infty}^\infty$. Let $E : \Omega \rightarrow \Omega$ be a function such that E^{-1} exists, and there exists a constant $C_E \in (0, \eta^{-1})$ satisfying the following Assumption (E1):

(E1) For each $n \in \mathbb{Z}$ one has

$$d(\sigma^n E^{-1}x, \sigma^n E^{-1}y) \leq C_E d(\sigma^n x, \sigma^n y).$$

Now define the coupled map $T : \Omega \rightarrow \Omega$ by $T(x) = E(\bar{\tau}(x))$. Define, for $\Phi \in \mathcal{C}$, $\mathcal{L}(\Phi)(x) = L(\Phi)(E^{-1}(x))$. It is straightforward to verify that for $\Phi, \Psi \in \mathcal{C}$, $\mathcal{L}(\Phi \circ T \cdot \Psi) = \Phi \cdot \mathcal{L}(\Psi)$ (as $T = E \circ \bar{\tau}$).

Theorem 4.1. *There exists a constant C_6 such that for every $\Phi \in \mathcal{C}$, $\mathcal{L}(\Phi) \in \mathcal{C}$ one has*

$$|\mathcal{L}^n(\Phi)|_\beta \leq |\Phi|_\beta (C_E \eta)^{\beta n} + C_6 |\Phi|_\infty C_E^\beta \sum_{i=0}^{\infty} (C_E \eta)^{i\beta}$$

Proof. By Lemma 3.10 ($n = 1$)

$$\begin{aligned} |\mathcal{L}(\Phi)(x) - \mathcal{L}(\Phi)(y)| &= |L(\Phi)(E^{-1}(x)) - L(\Phi)(E^{-1}(y))| \\ &\leq |L(\Phi)|_\beta d(E^{-1}x, E^{-1}y)^\beta \\ &\leq (|\Phi|_\beta \eta^\beta + C_5 |\Phi|_\infty) C_E^\beta d(x, y)^\beta \end{aligned}$$

and so

$$|\mathcal{L}(\Phi)|_\beta \leq |\Phi|_\beta (C_E \eta)^\beta + C_5 C_E^\beta |\Phi|_\infty.$$

Assume the formula for $n - 1$. Since $|\mathcal{L}(\Phi)|_\infty = |L(\Phi)|_\infty \leq |\Phi|_\infty$, we obtain inductively

$$\begin{aligned} |\mathcal{L}^n(\Phi)|_\beta &\leq |\mathcal{L}^{n-1}(\Phi)|_\beta (C_E \eta)^\beta + C_5 C_E^\beta |\Phi|_\infty \\ &\leq |\Phi|_\beta (C_E \eta)^{n\beta} + C_5 |\Phi|_\infty C_E^\beta \sum_{i=1}^{\infty} (C_E \eta)^{\beta i} + C_5 C_E^\beta |\Phi|_\infty \\ &= |\Phi|_\beta (C_E \eta)^{n\beta} + C_5 |\Phi|_\infty C_E^\beta \sum_{i=0}^{\infty} (C_E \eta)^{\beta i}. \end{aligned}$$

□

Let us note that $C_4 = C_5 = C_6$. We record, for future use, two elementary topological properties of the map $T : \Omega \rightarrow \Omega$.

Lemma 4.2. *For each $x \in \Omega$, $\{T^{-n}x, n \geq 0\}$ is dense in Ω .*

Proof. Let $x \in \Omega$, and let $\epsilon > 0$. Since $\{\tau^{-1}x_0\} \cup \{0, 1\}$ forms a partition of $[0, 1]$ with mesh $< \eta$, $\bar{\tau}^{-1}(E^{-1}x)|_{\Lambda_0}$ also forms a partition with mesh smaller than $C_E \eta < 1$. On iterating, we see that $T^{-n}(x)|_{\Lambda_0}$ generates a partition with mesh less than $(C_E \eta)^n$. This remains true on each node of the lattice; denoting by Δ_i the mesh of the i th node of the lattice and defining $\Delta = \sup_i \theta^{|i|} \Delta_i$, we see that there will be a pre-image of x within ϵ of an arbitrarily chosen point y provided $n \geq \log(1/\epsilon) / \log(1/C_E \eta)$. □

Lemma 4.3. *If U and V are any open subsets of Ω , then there exists an $N \in \mathbb{Z}^+$ such that for all $n \geq N$, $U \cap T^n V \neq \emptyset$.*

Proof. Let $x \in U$. Since $\{T^{-n}x, n \geq 0\}$ is dense in Ω , there exists an n_1 such that $T^{-n_1}x \in V$. Since V is open, there exists a $\delta > 0$ such that $B_\delta(T^{-n_1}x) \in V$. Choose $n_2 \geq \log(2/\delta) / \log(1/C_E \eta)$. Then for $n \geq n_2$, there always exists an element of $\{T^{-n}x\}$ inside $B_\delta(T^{-n_1}x)$ since the mesh Δ as defined above is less than δ for $n \geq n_2$. Hence $T^{-n}U \cap V \neq \emptyset \forall n \geq n_2$. □

We note that the above lemmas imply that the map T is topologically mixing, and since Ω is compact, forward transitive. Further, the map T is expansive, namely, if $x \in \Omega$ and $y \in \Omega$ are such that $T^n x = T^n y$ for every $n \in \mathbb{N}$ then $x = y$.

5. SPECTRAL PROPERTIES OF \mathcal{L} .

5.1. Ionescu-Tulcea and Marinescu Theorem. In this section we recall the classical machinery of quasi-compactness that is used to establish the statistical properties of a deterministic dynamical system, once a Lasota-Yorke type inequality (see Theorem 4.1) can be established. The main ingredient is the theorem by Ionescu-Tulcea and Marinescu.

Definition 5.1. *Let L be an operator on a Banach space $(V, \|\cdot\|)$. L is quasi-compact if there exists a positive integer r and a compact operator K such that $\|L^r - K\| < 1$.*

If L is quasi-compact, then $V = F \oplus H$ with F and H invariant under L , $\dim F < \infty$, $r(L|_H) < r(L)$ and each eigenvalue of $L|_F$ has modulus $r(L)$, where $r(\cdot)$ denotes the spectral radius.

As noted in [BG97, DZ09], quasi compactness can be restated in many equivalent ways. We give below the following definition [DZ09] as this is the form in which we will use quasi-compactness.

Definition 5.2 (Theorem 2.5.3, [DZ09]). *$L : V \rightarrow V$ is quasi-compact if and only if there are bounded linear operators $\{Q_\sigma : \sigma \in \Upsilon\}$ and R on V such that*

$$\begin{aligned} L^n &= \sum_{\sigma \in \Upsilon} \sigma^n \phi_\sigma + R^n \quad \forall \quad n = 1, 2, \dots, \\ \phi_\sigma \phi_{\sigma'} &= 0 \text{ if } \sigma \neq \sigma' \\ \phi_\sigma^2 &= \phi_\sigma, \quad \forall \quad \sigma \in \Upsilon \\ \phi_\sigma R &= R \phi_\sigma = 0 \quad \forall \quad \sigma \in \Upsilon \\ \phi_\sigma V &= D(\sigma), \quad \forall \quad \sigma \in \Upsilon \\ r(R) &< 1 \end{aligned}$$

where Υ is the set of the eigenvalues of L with modulus 1, $D(\sigma) = \{f \in V : Lf = \sigma f\}$ is the eigen-space of L corresponding to the eigenvalue σ and $r(R) := \lim_{n \rightarrow \infty} \|R^n\|^{1/n}$ is the spectral radius of R .

Next, we recall a version of the Ionescu-Tulcea and Marinescu theorem, established by Hennion and Hervé [HH01, Theorem II.5].

Theorem 5.3. *Let $|\cdot|$ be a continuous semi-norm on a Banach space $(V, \|\cdot\|)$ and let Q be a bounded operator on V such that*

(1)

$$Q(\{f : f \in V, \|f\| \leq 1\})$$

is conditionally compact in $(V, |\cdot|)$

(2) *there exists a constant M such that for all $f \in V$, $|Q(f)| \leq M|f|$*

(3) *there exists a $k \in \mathbb{N}$ and real numbers r and R such that $r < r(Q)$ and for all $f \in V$*

$$\|Q^k f\| \leq R|f| + r^k \|f\|.$$

Then Q is quasi-compact.

5.2. Quasi-compactness and other properties of \mathcal{L} , P .

Lemma 5.4. *\mathcal{L} is quasi-compact.*

Proof. We will take $|\cdot| = |\cdot|_\infty$ and $\|\cdot\| = |\cdot|_\beta + |\cdot|_\infty$. The set \mathcal{C} is a Banach space under $\|\cdot\|$. Clearly, condition 1 is satisfied since the unit ball of $(V, \|\cdot\|)$ is mapped under \mathcal{L} inside a ball of finite radius. Condition 2 is true because $|\mathcal{L}(\Phi)|_\infty \leq |\Phi|_\infty$. Condition 3 follows very easily from Theorem 4.1 and the observation that $\mathcal{L}\mathbf{1} = \mathbf{1}$ implies that $r(\mathcal{L}) \geq 1$. \square

The next sequence of lemmas establish that 1 is the unique eigenvalue of \mathcal{L} on the unit disk, and that 1 is a simple eigenvalue. The proofs of these statements follow standard lines (see, for instance, the proof of the Ruelle-Perron-Frobenius Theorem, [Bal00]), and require that the map T be forward transitive, and topologically mixing. We have to, however, account for the non-standard definition of \mathcal{L} .

As a technical point, we observe that the operators \mathcal{L}_k defined as

$$\mathcal{L}_k(\Phi)(x) = L_k(\Phi)(E^{-1}x)$$

are point-wise approximations to the operator \mathcal{L} , that is, $\lim_{k \rightarrow \infty} \mathcal{L}_k(\Phi)(x) = \mathcal{L}(\Phi)(x)$ for each $\Phi \in \mathcal{C}$ and $x \in \Omega$.

Lemma 5.5. *1 is a simple eigenvalue for \mathcal{L} with eigenfunction $\mathbf{1}$.*

Proof. First, we show that any eigenfunction ψ for \mathcal{L} to the eigenvalue 1 is either zero, or nowhere vanishing.

Since \mathcal{L} is a real linear operator it is enough to consider real valued eigenfunctions.

First we show that any real valued eigenfunction can be written as a linear combination of non-negative eigenfunctions. Let ψ^+ and ψ^- be the positive and negative parts of ψ . Since $\psi^\pm \leq |\psi|$, $\psi^\pm \in \mathcal{C}$. Further, the set

$$F_+ := \left\{ \frac{1}{n} \sum_{j=0}^{n-1} \mathcal{L}^j \psi^+ \mid n \geq 1 \right\}$$

has a bounded diameter in the Lipschitz norm and hence is equicontinuous and bounded in the $|\cdot|_\infty$ norm. The Arzelà-Ascoli theorem implies that there exists a subsequence n_j such that $\lim_{n_j \rightarrow \infty} \frac{1}{n_j} \sum_{j=0}^{n_j-1} \mathcal{L}^j(\psi^+) \rightarrow \psi_\infty^+$ uniformly in the $|\cdot|_\infty$ norm. Clearly, $\psi_\infty^+ \geq 0$, and $\mathcal{L}\psi_\infty^+ = \psi_\infty^+$. Finally, $|\psi_\infty^+|_\infty < \infty$ and by Theorem 4.1, $|\psi_\infty^+|_\beta < \infty$; this implies that $\psi_\infty^+ \in \mathcal{C}$. A similar analysis can be performed for ψ^- . It then follows from a straightforward diagonalization argument that

$$\psi = \lim_{n_j \rightarrow \infty} \frac{1}{n_j} \sum_{j=0}^{n_j-1} \mathcal{L}^j \psi^+ - \frac{1}{n_j} \sum_{j=0}^{n_j-1} \mathcal{L}^j \psi^- = \psi_\infty^+ - \psi_\infty^-.$$

For the rest of this proof, it will be assumed that all eigenfunctions are non-negative and real. Let $x \in \Omega$ be a point such that $\psi(x) = 0$, with ψ an eigenfunction. Then

$$0 = \psi(x) = \mathcal{L}^n(\psi)(x) = \lim_{k \rightarrow \infty} \mathcal{L}_k^n(\psi)(x) = \lim_{k \rightarrow \infty} \sum_{|\zeta|=k} \frac{1}{b_k} e^{g^{(n)}(i_k \zeta_x)} \psi(i_k \zeta_x).$$

By Lemma 4.2 and the non-negativity and continuity of ψ , it follows that ψ is identically 0.

To show that the geometric multiplicity of 1 is 1, suppose there are two positive real eigenfunctions ϕ and ψ and put

$$t = \inf_{x \in \Omega} \frac{\phi(x)}{\psi(x)},$$

which equals $\phi(z)/\psi(z)$ at some point $z \in \Omega$. Then $h(x) = \phi(x) - t\psi(x)$ is an eigenfunction to the eigenvalue 1, and $h(z) = 0$ which, by the previous paragraph, implies that $h \equiv 0$. Therefore ψ is some multiple of ϕ .

Finally, we show that the algebraic multiplicity of 1 is also 1. Suppose not. Then there exists a ψ with $(1 - \mathcal{L})^2 \psi = 0$ but $(1 - \mathcal{L})\psi \neq 0$. Since $(1 - \mathcal{L})\psi$ is an eigenvector for \mathcal{L} , we must have

$\mathcal{L}\psi - \psi = k\mathbf{1}$ for some $k \neq 0$. Iteration yields $\mathcal{L}^n\psi = nk\mathbf{1} + \psi$ which contradicts the uniform boundedness of \mathcal{L}^n as $|\mathcal{L}^n\psi|_\infty \leq |\psi|_\infty \forall n$. \square

Lemma 5.6. *Let $\mathcal{M}(\Omega)$ be the space of complex Radon measures on Ω . The operator $\mathcal{L}^* : \mathcal{M}(\Omega) \rightarrow \mathcal{M}(\Omega)$ is well defined. There exists a unique probability measure ν such that $\mathcal{L}^*(\nu) = \nu$.*

Proof. Note that \mathcal{C} is dense in $C(\Omega)$ (by the Stone-Weierstrass Theorem, since \mathcal{C} separates points, Ω is compact, Hausdorff, and \mathcal{C} contains the constant functions). Since \mathcal{L} is Lipschitz on \mathcal{C} , it has a continuous extension to $C(\Omega)$, that we also denote by \mathcal{L} . Since the dual of $C(\Omega)$ is $\mathcal{M}(\Omega)$, the operator $\mathcal{L}^* : \mathcal{M} \rightarrow \mathcal{M}$ is well defined. Since \mathcal{L} and \mathcal{L}^* have the same spectrum, 1 is also a simple eigenvalue for \mathcal{L}^* . By Lemma 5.5 we conclude that there exists a probability measure ν such that $\mathcal{L}^*\nu = \nu$.

To establish the uniqueness of ν we prove that 1 is the only eigenvalue for \mathcal{L} on the unit disk. Let λ be another eigenvalue of modulus 1, and let ϕ_λ be the eigenvector corresponding to λ . By orthogonality, $\int \phi_\lambda d\nu = 0$. But, for any $\psi \in \mathcal{C}$, we will show that $\mathcal{L}^n(\psi) \rightarrow \int \psi d\nu$. This will then give us a contradiction, because $\mathcal{L}^n\phi_\lambda = \lambda^n\phi_\lambda$ does not converge to 0.

We note that it is sufficient to prove this claim for positive $\psi \in \mathcal{C}$ because if $\psi' \in \mathcal{C}$, then on decomposing $\psi' = \psi^+ - \psi^-$ we get $\mathcal{L}^n\psi' = \mathcal{L}^n\psi^+ - \mathcal{L}^n\psi^- \rightarrow \int (\psi'^+ - \psi'^-) d\nu = \int \psi' d\nu$. Now, if ψ is some positive element of \mathcal{C} , denote by $\tilde{\psi}$ any continuous accumulation point of $\mathcal{L}^n\psi$. We will show that $\tilde{\psi}$ must be constant.

To see this, observe that $\tilde{\psi} \geq 0$ and

$$\sup \tilde{\psi} \geq \sup \mathcal{L}\tilde{\psi} \geq \dots \geq \sup \mathcal{L}^n\tilde{\psi} \geq \dots$$

Since $\tilde{\psi}$ is an accumulation point for $\mathcal{L}^n\psi$ none of these inequalities can be strict, so in fact $\sup \mathcal{L}^n\tilde{\psi} = \sup \tilde{\psi}$ for all $n \geq 0$. By continuity, there exists a point x_n with $\sup \mathcal{L}^n\tilde{\psi} = \mathcal{L}^n\tilde{\psi}(x_n)$. This implies that

$$\tilde{\psi}(x_0) = \mathcal{L}^n\tilde{\psi}(x_n) = \lim_{k \rightarrow \infty} \frac{1}{b_k} \sum_{|\zeta|=k} e^{g^{(n)}(i_k\zeta_{x_n})} \tilde{\psi}(i_k\zeta_{x_n})$$

from where it follows that

$$\lim_{k \rightarrow \infty} \left| \frac{1}{b_k} \sum_{|\zeta|=k} e^{g^{(n)}(i_k\zeta_{x_n})} \left(\tilde{\psi}(x_0) - \tilde{\psi}(i_k\zeta_{x_n}) \right) \right| = 0.$$

Using again the fact that

$$\frac{1}{b_k} \sum_{|\zeta|=k} e^{g^{(n)}(i_k\zeta_{x_n})} = \mathcal{L}_k^n \mathbf{1}(x_n) \rightarrow 1 \text{ as } k \rightarrow \infty$$

for each $n \geq 0$, we get that, on writing $\mathcal{L}_k^n \mathbf{1}(x_n) = 1 + \epsilon_k$ with $\epsilon_k \rightarrow 0$, and observing that $\tilde{\psi}(x_0) \geq \tilde{\psi}(i_k\zeta)$ for each branch,

$$\lim_{k \rightarrow \infty} (1 + \epsilon_k)(\tilde{\psi}(x_0) - \tilde{\psi}(i_k\zeta_{x_n})) = 0.$$

Since for each branch $\lim_{k \rightarrow \infty} i_k\zeta_{x_n} = y$ where y satisfies $T^n y = x_n$ we have that $\tilde{\psi}(y) = \tilde{\psi}(x_0)$ for all $y \in \{T^{-n}x_n\}$ for each $n \geq 0$. It follows from lemma 4.3 that $\cup_n \{T^{-n}x_n\}$ is dense, and so $\tilde{\psi}$ is constant.

Therefore,

$$\tilde{\psi} = \int \tilde{\psi} d\nu = \int \mathcal{L}^n \tilde{\psi} d\nu = \int \psi d\nu$$

where the last equality uses the fact that $\mathcal{L}^*\nu = \nu$. \square

We now show that the measure ν constructed above is conformal. Recall that for any measurable function f , a measure μ is said to be f -conformal if for every measurable set A on which the map $T : A \rightarrow TA$ is invertible,

$$(3) \quad \int_A e^{-f} d\mu = \mu(TA).$$

In the context of continuous transformations T on a compact metric space Ω , when the map T has a finite generating partition, one requires that (3) hold for all measurable A . In our setup, we do not obtain a finite, or countable, generating partition for the map T . Therefore, we only check (3) on open sets.

Proposition 5.7. *The measure ν is g -conformal, i.e., if A be an open set on which $T : A \rightarrow TA$ is invertible, then*

$$\int_A e^{-g} d\nu = \nu(TA).$$

Proof.

$$\begin{aligned} \int_A e^{-g} d\nu &= \int_{\Omega} \mathcal{L}(\chi_A e^{-g})(x) d\nu(x) \\ &= \int_{\Omega} \lim_{k \rightarrow \infty} (1/b_k) \sum_{\zeta \in T^{-1}(\pi_k x)} e^{g(i_k \zeta)} e^{-g(i_k \zeta)} \chi_A(i_k \zeta) d\nu(x) \\ &= \int_{\Omega} \lim_{k \rightarrow \infty} (1/b_k) \sum_{\zeta \in T^{-1}(\pi_k x)} \chi_A(i_k \zeta) d\nu(x) \\ &= \int_{\Omega} \lim_{k \rightarrow \infty} (1/b_k) \chi_{TA}(x) b_k d\nu(x) \\ &= \nu(TA). \end{aligned}$$

The third equality from the bottom follows by observing that $\zeta \in T^{-1}(x) \in A$ if and only if $x \in TA$. \square

Theorem 5.8. *There exists a constant $0 < \varsigma < 1$ with the property that for any $\phi_1, \phi_2 \in \mathcal{C}$ there exists a constant C_7 such that*

$$\left| \int \phi_1 \circ T^n \phi_2 d\nu - \int \phi_1 d\nu \int \phi_2 d\nu \right| \leq C_7 |\phi_1|_{\infty} \|\phi_2\| \varsigma^n.$$

Proof. \mathcal{L} has a unique eigenvalue of modulus 1, therefore the projection operator on the eigen-space of 1 is defined by $\mathcal{P}(\phi) = \int \phi d\nu$. Since \mathcal{L} is quasi-compact, there must exist a constant $\varsigma \in (0, 1)$ such that (for some c_1)

$$\|\mathcal{L}^n(\phi - \mathcal{P}(\phi))\| \leq c_1 \varsigma^n \|\phi\|.$$

Let $\tilde{\phi}_1 = \phi_1 - \int \phi_1 d\nu$. Observe that

$$\left| \int \phi_1 \circ T^n \phi_2 d\nu - \int \phi_1 d\nu \int \phi_2 d\nu \right| = \left| \int \tilde{\phi}_1 \circ T^n \phi_2 d\nu \right| = \left| \int \tilde{\phi}_1 \mathcal{L}^n \phi_2 d\nu \right|$$

which is bounded by $|\tilde{\phi}_1|_{\infty} c_1 \varsigma^n \|\phi_2\| + \int \tilde{\phi}_1 d\nu \int \mathcal{L}^n \mathcal{P} \phi_2 d\nu$ with the second term being 0. On adjusting the constant c_1 , we finish the proof. \square

6. ALMOST SURE INVARIANCE PRINCIPLE

Theorem 6.1. *The invariant measure ν satisfies the almost sure invariance principle.*

For a function $f \in \mathcal{C}$ (note that this $f \in L^\infty(\nu)$) and for t sufficiently close to 0, define a family of operators $\mathcal{L}_{t,f}, t \in \mathbb{R}$, on \mathcal{C} by

$$\mathcal{L}_{t,f}(\Phi) = \mathcal{L}\left(e^{itf}\Phi\right).$$

We now recall an abstract theorem by Gouëzel [Gou10], stated in a manner relevant to our setting.

Proposition 6.2. *Suppose \mathcal{L} satisfies Theorem 4.1 and there exists a constant $C_8 > 0$ such that $\|\mathcal{L}_{f,t}^n\|_{\mathcal{C} \rightarrow \mathcal{C}} \leq C_8$ for all $n \in \mathbb{N}$ and for all t small enough. Then there exists a probability space (Γ) and two processes (A_j) and (B_j) on Γ such that*

- (1) *the processes $(f \circ T^j)$ and (A_j) have the same distribution*
- (2) *the random variables (B_j) are independent and distributed as $\mathcal{N}(0, \sigma^2)$ for an appropriately chosen σ^2 , and*
- (3) *almost surely in Γ*

$$\left| \sum_{l=0}^{n-1} A_j - \sum_{l=0}^{n-1} B_j \right| = o(n^\gamma)$$

for any $\gamma > 0.25$.

As noted in their paper, since a Brownian motion at integer times coincides with a sum of iid Gaussian random variables, this theorem can be formulated as an almost sure approximation by a Brownian motion. To establish the almost sure invariance principle for our setup, we only need to check that $\|\mathcal{L}_{f,t}^n\|_{\mathcal{C} \rightarrow \mathcal{C}}$ stays bounded for t small enough.

Lemma 6.3. *There exists a constant $C_9 > 0$ such that $\|\mathcal{L}_{f,t}^n\|_{\mathcal{C} \rightarrow \mathcal{C}} \leq C_9$ for all $n \in \mathbb{N}$ and for all t small enough.*

Proof. Since $|\mathcal{L}_{f,t}^n(\Phi)|_\infty \leq |e^{itf}\Phi|_\infty \leq |\Phi|_\infty$, we only need to check if $|\mathcal{L}_{f,t}^n(\Phi)|_\beta$ is bounded. By Theorem 4.1

$$|\mathcal{L}_{f,t}^n(\Phi)|_\beta \leq |e^{itf}\Phi|_\beta (C_E \eta)^n + C_6 |\Phi|_\infty.$$

It remains to bound $|e^{itf}\Phi|_\beta$. Indeed, by the triangle inequality:

$$\begin{aligned} \left| e^{itf(x)}\Phi(x) - e^{itf(y)}\Phi(y) \right| &\leq \left| e^{itf(x)}\Phi(x) - e^{itf(x)}\Phi(y) \right| + \left| e^{itf(x)}\Phi(y) - e^{itf(y)}\Phi(y) \right| \\ &\leq |\Phi(x) - \Phi(y)| + |\Phi|_\infty |e^{itf(x)} - e^{itf(y)}| \\ &\leq |\Phi|_\beta d(x, y)^\beta + |\Phi|_\infty c_1 d(x, y)^\beta + o(d(x, y)^\beta) \end{aligned}$$

where the constant c_1 depends on f (as well as η and β). The last inequality uses an argument from lemma 3.9. On dividing by $d(x, y)^\beta$ and taking the supremum we obtain (for some c_2)

$$|e^{itf}\Phi|_\beta \leq |\Phi|_\beta + c_2 |\Phi|_\infty.$$

Hence we can put

$$C_9 := \max \left\{ \sup_n \{ (|\Phi|_\beta + c_2 |\Phi|_\infty) (C_E \eta)^n \} + C_6, 1 \right\}$$

which is finite. □

REFERENCES

- [Bal00] Viviane Baladi. *Positive transfer operators and decay of correlations*, volume 16 of *Advanced Series in Nonlinear Dynamics*. World Scientific Publishing Co. Inc., River Edge, NJ, 2000.
- [Bar02] Jean-Baptiste Bardet. Limit theorems for coupled analytic maps. *Probab. Theory Related Fields*, 124(2):151–177, 2002.
- [BC91] Michael Benedicks and Lennart Carleson. The dynamics of the Hénon map. *Ann. of Math. (2)*, 133(1):73–169, 1991.
- [BG97] Abraham Boyarsky and Paweł Góra. *Laws of chaos*. Probability and its Applications. Birkhäuser Boston Inc., Boston, MA, 1997. Invariant measures and dynamical systems in one dimension.
- [BGK07] Jean-Baptiste Bardet, Sébastien Gouëzel, and Gerhard Keller. Limit theorems for coupled interval maps. *Stoch. Dyn.*, 7(1):17–36, 2007.
- [BLMMR92] L. A. Bunimovich, R. Livi, G. Martínez-Mekler, and S. Ruffo. Coupled trivial maps. *Chaos*, 2(3):283–291, 1992.
- [Bow08] Rufus Bowen. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, volume 470 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, revised edition, 2008. With a preface by David Ruelle, Edited by Jean-René Chazottes.
- [BS88] L. A. Bunimovich and Ya. G. Sinai. Spacetime chaos in coupled map lattices. *Nonlinearity*, 1(4):491–516, 1988.
- [Bun97] Leonid A. Bunimovich. Coupled map lattices: some topological and ergodic properties. *Phys. D*, 103(1-4):1–17, 1997. Lattice dynamics (Paris, 1995).
- [BY93] Michael Benedicks and Lai-Sang Young. Sinai-Bowen-Ruelle measures for certain Hénon maps. *Invent. Math.*, 112(3):541–576, 1993.
- [DZ09] Jiu Ding and Aihui Zhou. *Statistical properties of deterministic systems*. Tsinghua University Texts. Springer-Verlag, Berlin, 2009.
- [Gou10] Sébastien Gouëzel. Almost sure invariance principle for dynamical systems by spectral methods. *Annals of Probability*, 38(4):1639–1671, 2010.
- [GR93a] V. M. Gundlach and D. A. Rand. Spatio-temporal chaos. I. Hyperbolicity, structural stability, spatio-temporal shadowing and symbolic dynamics. *Nonlinearity*, 6(2):165–200, 1993.
- [GR93b] V. M. Gundlach and D. A. Rand. Spatio-temporal chaos. II. Unique Gibbs states for higher-dimensional symbolic systems. *Nonlinearity*, 6(2):201–213, 1993.
- [GR93c] V. M. Gundlach and D. A. Rand. Spatio-temporal chaos. III. Natural spatio-temporal measures for coupled circle map lattices. *Nonlinearity*, 6(2):215–230, 1993.
- [HH01] Hubert Hennion and Loïc Hervé. *Limit theorems for Markov chains and stochastic properties of dynamical systems by quasi-compactness*, volume 1766 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001.
- [KK92] Gerhard Keller and Martin Künzle. Transfer operators for coupled map lattices. *Ergodic Theory Dynam. Systems*, 12(2):297–318, 1992.
- [KKN92] Gerhard Keller, Martin Künzle, and Tomasz Nowicki. Some phase transitions in coupled map lattices. *Phys. D*, 59(1-3):39–51, 1992.
- [KL04] Gerhard Keller and Carlangelo Liverani. Coupled map lattices without cluster expansion. *Discrete Contin. Dyn. Syst.*, 11(2-3):325–335, 2004.
- [KL05] G. Keller and C. Liverani. A spectral gap for a one-dimensional lattice of coupled piecewise expanding interval maps. In *Dynamics of coupled map lattices and of related spatially extended systems*, volume 671 of *Lecture Notes in Phys.*, pages 115–151. Springer, Berlin, 2005.
- [KL06] Gerhard Keller and Carlangelo Liverani. Uniqueness of the SRB measure for piecewise expanding weakly coupled map lattices in any dimension. *Comm. Math. Phys.*, 262(1):33–50, 2006.
- [PT92] Antonio Politi and Alessandro Torcini. Periodic orbits in coupled Hénon maps: Lyapunov and multifractal analysis. *Chaos*, 2(3):293–300, 1992.
- [Rug02] Hans Henrik Rugh. Coupled maps and analytic function spaces. *Ann. Sci. École Norm. Sup. (4)*, 35(4):489–535, 2002.
- [WY08] Quidong Wang and Lai-Sang Young. Toward a theory of rank one attractors. *Ann. of Math. (2)*, 167(2):349–480, 2008.
- [You02] Lai-Sang Young. What are SRB measures, and which dynamical systems have them? *J. Statist. Phys.*, 108(5-6):733–754, 2002. Dedicated to David Ruelle and Yasha Sinai on the occasion of their 65th birthdays.

(Chinmaya Gupta) UNIVERSITY OF SOUTHERN CALIFORNIA, 3620 S. VERMONT AVE, LOS ANGELES, CA, 90089
E-mail address: chinmaya.gupta@usc.edu
URL: <http://www-bcf.usc.edu/~chinmayg>

(Nicolai Haydn) UNIVERSITY OF SOUTHERN CALIFORNIA, 3620 S. VERMONT AVE, LOS ANGELES, CA, 90089
E-mail address: nhaydn@usc.edu